

Force Free Waves and Black Hole Magnetospheric Causality

Brian Punsly

4014 Emerald Street No.116, Torrance CA, USA 90503

brian.punsly@hsc.com or brian.punsly@gte.net

ABSTRACT

The force free approximation is often useful when describing tenuous plasmas in strong cosmic magnetic fields. Time evolution of any such system is governed by the information that can be transported along the characteristics of the plasma modes allowed by the force free constraint. This brief article elucidates, for the first time, the nature of the information that can be transported along the characteristics for each of the two plasma modes, the Alfvén and fast modes. It is reassuring that these modes behave as one would expect if the perfect magnetohydrodynamic (MHD) modes in a magnetically dominated plasma were evaluated in the limit of zero plasma inertia (i.e., only allowed to propagate information consistent with the force free constraint). There are no properties of these waves that do not exist within the MHD theory, when this additional constraint on the current is imposed. The results of the characteristic analysis elucidates the nature of the causality violation in force free black hole magnetospheres. The most significant result from the standpoint of global causality is that charge and current perturbations can not be transported along the fast mode characteristics in the force free limit.

1. Introduction

Applications of force free electrodynamics are common in the astrophysical literature. The assumption that the plasma inertia is negligible compared to the magnetic energy density is an accurate depiction of the plasma state in many contexts. This condition justifies the force free approximation based on a local physical frame analysis. The most notable successes of such an approach are descriptions of equilibria in the solar magnetosphere. Motivated by the solar work, it has also been extended to studies of the physics of black hole magnetospheres (see Blandford and Znajek (1977) and Thorne et al (1986)) and more recently gamma ray bursts (see Blandford (2002)).

However, the implementation of the force free condition to black hole magnetospheres is not so trivial because general relativity is not a local theory. In particular, the concepts of an event horizon and ergosphere are globally defined (Hawking 1973). Black hole magnetospheres are the main focus of this treatment, **yet all of the results that are derived in this article apply equally well to any force free calculation.** A salient example of the global nature of general relativity is the fact the physical boundary condition at the event horizon is that all particles that are near the black hole must corotate with the event horizon and go into the black hole on trajectories that approach the ingoing light-like principal null congruence, as seen by any external observer (see Chapter 3 of Punsly (2001)). This applies to any plasma, no matter how tenuous and is independent of the magnetic field strength. There are no stable equilibria near the horizon, gravity always overwhelms the other forces. This property is precisely what distinguishes a black hole from another type of compact object such as a neutron star or white dwarf. All plasma near the horizon is inertially dominated: a global condition. In a sense, the plasma inertia is infinite near the horizon as viewed globally. By contrast, a local analysis would lead one to believe that the force free approximation is valid (i.e., the plasma inertia is negligible). In spite of this defining property of the black hole, force free treatments of black hole magnetospheres that extend all the way to the event horizon have been proposed by many authors (most notably Blandford and Znajek (1977) and Thorne et al (1986)).

In this paper, we consider the plasma waves in a force free magnetosphere. This is important in order to clarify certain anecdotal defenses of the force free approximation near the event horizon which infer that force free plasma waves can transport different information than their counterparts in the full perfect magnetohydrodynamic (MHD) treatment of plasma waves in magnetically dominated magnetospheres (see Chapter 2 of Punsly (2001) for a detailed MHD treatment and Blandford (2001) for the force free discussion). In this article, it is shown by means of a straightforward explicit calculation that this claim is erroneous. This article elucidates the current structure within the individual wave modes for the first time (even the lengthy, complicated and very detailed analysis of certain topics of Uchida (1997) in the context of black hole magnetospheres does not address these most relevant issues in a force free magnetosphere).

2. The Force Free Approximation and Magnetically Dominated MHD

This section compares and contrasts the force free approximation with a magnetically dominated perfect MHD system. Generally, the force free approximation is considered as a zero mass limit of the full MHD theory. It is usually considered an expedience that reduces

the dimension of the full MHD system of hyperbolic differential equations to five. It is very rarely argued to be a more accurate physical description than the full MHD theory (but, see Blandford (2002) for such an argument). Generally, the force free approximation is invoked is to simplify the mathematical treatment not to improve on the physical depiction. If the force free approximation is not pathological then one would expect it to occur as the natural limit of zero inertia in the full MHD theory. This section begins to lay the groundwork to show that this is indeed the case. The set of force free equations are the current constraint written covariantly in terms of the Maxwell tensor and in a local inertial coordinate system,

$$F^{\mu\nu} J_\nu = 0 , \quad (2-1a)$$

$$\rho_e \mathbf{E} + \frac{\mathbf{J} \times \mathbf{B}}{c} = 0 , \quad (2-1b)$$

and Maxwell's equations (written in local coordinates)

$$\nabla \times \mathbf{B} = \frac{1}{c} \frac{\partial \mathbf{E}}{\partial t} + \frac{4\pi}{c} \mathbf{J} , \quad (2-2)$$

$$\nabla \times \mathbf{E} = -\frac{1}{c} \frac{\partial \mathbf{B}}{\partial t} , \quad (2-3)$$

$$\nabla \cdot \mathbf{B} = 0 , \quad (2-4)$$

$$\nabla \cdot \mathbf{E} = 4\pi\rho_e . \quad (2-5)$$

Equation (2.1) also implies the conditions:

$$\mathbf{E} \cdot \mathbf{B} = 0 , \quad (2-6)$$

$$\mathbf{E} \cdot \mathbf{J} = 0 . \quad (2-7)$$

The force free condition in equation (2.1) arises from the zero mass density limit of perfect MHD, the plasma has no inertia to stress the magnetic field lines.

3. Force Free Plasma Waves

The causal structure of a force free system is governed by the waves supported in the force free plasma. In this section we derive the nature of the information that can be transported along each of the wave characteristics. Recall that in perfect MHD the Alfven mode can transport charge and current that was either field aligned (force free) or cross-field (inertial). The fast wave cannot transport charge or field aligned currents, only inertial currents (see Punsly (2001), Chapter 2). It turns out that the effects of the force free constraint are equivalent to the zero mass density limit of perfect MHD plasma waves, as expected. Since the fast wave still cannot transport force free current in the zero inertia

limit, when the force free constraint on the current is imposed, the only value of current consistent with these two concepts (effectively two mathematical equations) is zero current. The fast mode in the force free limit carries no current or charge.

As a consequence of equation (2.6), one can find a frame in which the electric field vanishes if the Lorentz invariant $F^{\mu\nu}F_{\mu\nu} > 0$ (i.e., the field is magnetic). There exists a coordinate transformation to this frame that resembles a Lorentz boost in the azimuthal direction in an axisymmetric magnetosphere except that the boost parameter can exceed the speed of light. Thus, this is not a Lorentz transformation in general. However, the angular velocity as observed from asymptotic infinity in an asymptotically spherical coordinate system is referred to as the field line angular velocity, Ω_F , even when the azimuthal velocity is faster than the speed of light (Blandford 2002; Punsly 2001). Thus, this frame is often called the corotating frame of the magnetic field. The light cylinder is defined as the surface at which the azimuthal boost velocity equals the speed of light (in all physical frames). The implementation of this frame facilitates the calculations and one can perform a general coordinate transformation in order to evaluate plasma quantities in other physical frames. Since, this is not a Lorentz transformation in general, it is best to find coordinate invariant expressions. Another frame in which the electric field vanishes is the plasma rest frame. However, this frame is poorly defined in the force free limit since there is no accounting of the plasma state. Frames in which the electric field vanishes are sometimes called proper frames.

The calculations are performed in the short wavelength approximation. This is equivalent to the analysis found in Blandford (2002). No matter what the radius of curvature of the magnetic field, one can find plasma mode wavelengths that are short compared to this value. This is not very restrictive, even close to the event horizon of a black hole. One should be able to find proper frames in which the radius of curvature of the magnetic field is on the order of the geometrized mass of a black hole (10^{14} cm for a supermassive black hole). The unperturbed value of the four current, J_0^μ , effectively vanishes in the short wavelength approximation that the waves vary on much smaller scales than the background magnetic fields (the currents are effectively created by the gradients of the magnetic field). As a result of Ampere's law, this approximation is equivalent to treating the magnetic field of the unperturbed state as approximately a constant in the proper frame,

$$\frac{\partial}{\partial x^\alpha} \mathbf{B} \approx 0. \quad (3-1)$$

We analyze short wavelength linear (small amplitude) perturbations of equations (2.1)-(2.7). The waves are chosen in a plane wave representation to have an oscillatory behavior, $\exp i(\mathbf{k} \cdot \mathbf{r} - \omega t)$. We define the x direction to be parallel to $\mathbf{k}$, so all perturbed quantities

are Fourier analyzed in (x, t) space. For example, the four current is

$$J^\mu = J_0^\mu + e^{i(kx - \omega t)} \delta J^\mu \quad (3-2a)$$

$$\approx e^{i(kx - \omega t)} \delta J^\mu . \quad (3-2b)$$

The interested reader can find a discussion of the effects of field line curvature in a black hole magnetosphere in Chapter 6 of Punsly (2001). The nature of the resultant hybrid modes is beyond the scope of this treatment that has been chosen to parallel the analysis of Blandford (2002). In summary, field line curvature mixes the properties of the plasma modes as it does in the earth's magnetosphere (Lanzerotti and Southwood 1979). Long wavelength modes are neither fast nor Alfvén modes, but have hybrid properties (Appendix A is a derivation of the first order effects of magnetic field line curvature in the force free limit). The faster mode propagates slower as the wavelength increases and is no longer related to the critical point structure of the winds that reflect short wavelength mode speeds. The implication for black hole magnetospheres is that long wavelength force free fast modes travel slower than the speed of light and therefore can not propagate information from near the event horizon to asymptotic infinity.

The analysis of long wavelength plasma modes is very complicated and one is referred to the in depth analysis in Chapter 6 of Punsly (2001) to see the details of outgoing mode propagation resulting from field line curvature and spacetime curvature near the event horizon of a black hole. The results of those calculations are in accord with the causality arguments in this article. In particular, one finds that long wavelength plasma modes are extremely inefficient in their ability to transport magnetic stresses, field aligned current and charge outward from the spacetime near an event horizon.

In a frame with a vanishing electric field, the perturbed equations (2.1) - (2.7), with the condition above become:

$$\mathbf{B} \times \delta \mathbf{J} = 0 , \quad (3-3)$$

$$\omega \delta \mathbf{E} = -c \mathbf{k} \times \delta \mathbf{B} - i 4\pi \delta \mathbf{J} , \quad (3-4)$$

$$\omega \delta \mathbf{B} = c \mathbf{k} \times \delta \mathbf{E} , \quad (3-5)$$

$$\mathbf{k} \cdot \delta \mathbf{B} = 0 , \quad (3-6)$$

$$i \mathbf{k} \cdot \delta \mathbf{E} = 4\pi \delta \rho_e , \quad (3-7)$$

$$\delta \mathbf{E} \cdot \mathbf{B} = 0 . \quad (3-8)$$

It is useful to combine (2.2) and (2.3) to get the second order Maxwell's equation,

$$\nabla^2 \mathbf{E} - \nabla(\nabla \cdot \mathbf{E}) = \frac{1}{c^2} \frac{\partial^2 \mathbf{E}}{\partial t^2} + \frac{4\pi}{c^2} \frac{\partial \mathbf{J}}{\partial t} . \quad (3-9)$$

This equation gives the following relation amongst perturbed electromagnetic quantities,

$$k^2 \delta \mathbf{E} - \mathbf{k} (\mathbf{k} \cdot \delta \mathbf{E}) = \frac{\omega^2}{c^2} \delta \mathbf{E} + \frac{i4\pi\omega}{c^2} \delta \mathbf{J} . \quad (3-10)$$

At this point we explicitly write out the components of the perturbed equation above in a local Cartesian basis. The x-axis was previously defined to be parallel to $\mathbf{k}$ and we define the z-axis such that $\mathbf{B}$ always lies in the x-z plane. The components of the second order Maxwell's equation become:

$$\delta E_y = \frac{4\pi i\omega}{(c^2 k^2 - \omega^2)} \delta J_y , \quad (3-11a)$$

$$\delta E_z = \frac{4\pi i\omega}{(c^2 k^2 - \omega^2)} \delta J_z , \quad (3-11b)$$

$$\delta E_x = -\frac{4\pi i}{\omega} \delta J_x . \quad (3-11c)$$

Similarly an expansion of the components of the perturbed force free condition in (3.2) yields the following two conditions on the current:

$$\delta J_y = 0 , \quad (3-12)$$

$$\delta J_z = \tan \theta \delta J_x , \quad (3-13)$$

where θ is the angle between the propagation vector and the magnetic field direction. There is also a third constraint on the current that is obtained by substituting the perturbed second order Maxwell's equation (3.11) into the perturbed degeneracy condition (3.8):

$$\delta J_x (c^2 k^2 - \omega^2) = \omega^2 \frac{k_\perp}{k_\parallel} \delta J_z , \quad (3-14)$$

where $k_\parallel$ and $k_\perp$ are the components of the propagation vector parallel and perpendicular to the magnetic field direction, respectively.

Equation (3.14) can be substituted into the relation (3.13) in order to eliminate δJ_x . This yields the dispersion relation for plasma waves in a force free medium,

$$[\omega^2 - c^2 k^2] [\omega^2 - c^2 k_\parallel^2] = 0 . \quad (3-15)$$

3.1. The Fast Mode

The dispersion relation (3.15) has two solutions for the phase velocity. The fast mode is characterized by the phase velocity,

$$\frac{\omega}{k} = \pm c . \quad (3-16)$$

The group velocity is

$$\frac{\partial \omega}{\partial \mathbf{k}} = \pm c \frac{\mathbf{k}}{k} . \quad (3-17)$$

Inserting the value of the phase velocity into the perturbed induction equation, (3.5), and the perturbed degeneracy condition, (3.8), along with the vanishing divergence of the magnetic field, (3.6), and the constraint on the perturbed current, (3.14), we can solve for the electromagnetic components of the fast wave:

$$\delta B_x = 0 , \quad (3-18a)$$

$$\delta B_y = 0 , \quad (3-18b)$$

$$\delta B_z = \delta E_y , \quad (3-18c)$$

$$\delta E_x = 0 , \quad (3-18d)$$

$$\delta E_y = \delta B_z , \quad (3-18e)$$

$$\delta E_z = 0 . \quad (3-18f)$$

Substituting the values of the electric field from (3.18d-f) into the second order Maxwell's equation, (3.11), and the perturbed Gauss' law, (3.7), the light-like phase velocity of the fast wave requires that **all of the components of the four-current density that are propagated along the characteristics of the force free fast wave vanish identically**,

$$\delta J^\mu = 0 . \quad (3-19)$$

This is a coordinate invariant expression and is therefore true in all coordinate systems. The fast mode is linear in the force free limit, since there is a unique wave propagation speed given by (3.17). Thus, fast wave packets are linear superpositions of the oscillatory solutions. Therefore, the four-current density vanishes for fast wave packets as well.

From (3.18), one also has the following coordinate independent statements:

$$*\delta F^{\mu\nu} \delta F_{\mu\nu} = 0 , \quad (3-20a)$$

$$\delta F^{\mu\nu} \delta F_{\mu\nu} = 0 , \quad (3-20b)$$

$$*\delta F^{\mu\nu} F_{\mu\nu} = 0 , \quad (3-20c)$$

$$\delta F_{\mu\nu} U_G^\nu = 0 , \quad (3-20d)$$

$$\frac{\omega}{c} U_G^\nu = k^\nu , \quad (3-20e)$$

where U_G^ν is the group four velocity of the wave, which is expressed in the coordinates of the proper frame as:

$$U_G^\nu \equiv \left(c, \frac{\partial \omega}{\partial \mathbf{k}} \right) . \quad (3-21)$$

Not all of the relations in (3.20) are independent, for example (3.20d) implies (3.20a).

The primary result of this analysis, (3.19), that demonstrates the vanishing of the current in a fast mode is not a result of a biased interpretation by this author, a miscalculation or an unfounded assumption. The same result actually follows from the results of Blandford (2002). First of all, the wave structure that is described in a coordinate independent manner in (3.20) can be rewritten in the “laboratory frame coordinates” used by Blandford (2002). One then finds that, $\mathbf{k}$, $\delta\mathbf{B}$ and $\delta\mathbf{E}$ form an orthogonal triad as well as equations (7) and (8) of Blandford (2002). These are the results that are used in that treatment to describe the causal relevance of a fast wave to a force free magnetosphere. Inserting (7) and (8) into Ampere’s law, either (2.2) or (3.4), yields the fact that $\delta J_y = 0$ and $\delta J_z = 0$. By the orthogonality of the triad, $\delta E_x = 0$, as a consequence of the definition of the propagation direction. Then Ampere’s law can be used again to show that $\delta J_x = 0$. Finally, the vanishing of the charge density follows from either Gauss’ law or current conservation. Thus, (3.19) can be found from the analysis of Blandford (2002) quite readily.

3.2. The Alfven Wave

The other root of the dispersion relation (3.15) gives the phase velocity of the of the Alfven wave:

$$\frac{\omega}{k} = \pm c \cos \theta . \quad (3-22)$$

One can also compute the group velocity,

$$\frac{\partial \omega}{\partial \mathbf{k}} = \pm c \frac{\mathbf{B}}{B} . \quad (3-23)$$

Thus, the Alfven mode also propagates information at the speed of light, but only along the direction of the magnetic field.

Combining, the vanishing divergence of the perturbed magnetic field, (3.6), along with the constraint on the perturbed current, (3.14), and the induction equation, (3.5), inserted into the second order Maxwell’s equation, (3.11), one can solve for the components of the perturbed electromagnetic field transported along the characteristics of the Alfven mode:

$$\delta B_x = 0 , \quad (3-24a)$$

$$\delta B_y = -i \frac{4\pi k}{ck_{\perp}^2} \delta J_z = -i \frac{4\pi}{ck_{\perp}} \delta J_{\parallel} , \quad (3-24b)$$

$$\delta B_z = 0 , \quad (3-24c)$$

$$\delta E_x = -i \frac{4\pi}{\omega} \delta J_x = -i \frac{4\pi}{ck} \delta J_{\parallel} , \quad (3-24d)$$

$$\delta E_y = 0 , \quad (3-24e)$$

$$\delta E_z = -\cos \theta \delta B_y = -i \frac{4\pi k_{\parallel}}{ckk_{\perp}} \delta J_{\parallel} . \quad (3-24f)$$

The component of the current along the magnetic field in the proper frame is defined as

$$J_{\parallel} = \mathbf{J} \cdot \frac{\mathbf{B}}{B} . \quad (3-25)$$

The perturbed current can be found from (3.22), (3.24) and (3.14),

$$\delta J_x = i \frac{ck_{\perp} k_{\parallel}}{4\pi k} \delta B_y = \frac{k_{\parallel}}{k} J_{\parallel} , \quad (3-26a)$$

$$\delta J_y = 0 , \quad (3-26b)$$

$$\delta J_z = i \frac{ck_{\perp}^2}{4\pi k} \delta B_y = \frac{k_{\perp}}{k} J_{\parallel} , \quad (3-26c)$$

$$\delta \mathbf{J}_{\parallel} = \delta \mathbf{J} = i \frac{ck_{\perp}}{4\pi} \delta B_y \frac{\mathbf{B}}{B} . \quad (3-26d)$$

From Gauss' law, (3.7), and (3.24) the perturbed charge density is

$$\delta \rho_e = i \frac{k_{\perp}}{4\pi} \delta B_y = \frac{J_{\parallel}}{c} . \quad (3-27)$$

The second equality in (3.27) utilized the perturbed law of current conservation which is obtained from the combination of (3.7) and (3.4). The equations (3.24)-(3.26) can be combined to give the following useful relations amongst dynamical quantities that are propagated along the Alfven characteristics:

$$\delta \mathbf{J} \times \mathbf{B} = 0 , \quad (3-28a)$$

$$\delta \mathbf{E} \cdot \mathbf{B} = 0 , \quad (3-28b)$$

$$\delta \mathbf{E} \cdot \delta \mathbf{B} = 0 , \quad (3-28c)$$

$$\delta \mathbf{B} \cdot \mathbf{B} = 0 . \quad (3-28d)$$

Covariantly speaking, (3.28) implies the coordinate independent relations:

$$\delta J^{\mu} F_{\mu\nu} = 0 , \quad (3-29a)$$

$$*\delta F^{\mu\nu} F_{\mu\nu} = 0 , \quad (3-29b)$$

$$*\delta F^{\mu\nu} \delta F_{\mu\nu} = 0 , \quad (3-29c)$$

$$\delta F^{\mu\nu} F_{\mu\nu} = 0 , \quad (3-29d)$$

$$\delta F^{\mu\nu} \delta F_{\mu\nu} = 0 , \quad (3-29e)$$

$$\delta F_{\mu\nu} U_G^{\nu} = 0 , \quad (3-29f)$$

$$F_{\mu\nu} U_G^{\nu} = 0 . \quad (3-29g)$$

Equations (3.29a-d) are the coordinate independent analogs of the local expressions (3.28a-d). From (3.29a), the current flows along the magnetic field lines in a proper frame (this expression incorporates the assumption, (3.2), that field line curvature radii are large compared to the plasma mode wavelength). One can also get the orientation of the propagation vector relative to the perturbed and unperturbed fields in all frames by expressing (3.29a) in terms of the perturbed covariant version of Ampere's Law,

$$\delta F^{\mu\alpha} k_\alpha F_{\mu\nu} = 0 . \quad (3-30)$$

The Alfven mode transports a charge and a field aligned current (covariantly speaking, a force free current).

4. Causality in Force Free Magnetospheres

The calculation in the last section allows one to compute the characteristics of short wavelength waves in a force free magnetosphere. There are four modes: both ingoing and outgoing fast and Alfven waves. The characteristics by themselves are not enough to time evolve a force free system, one must also know what information can be propagated along these curves. The force free waves carry only two independent pieces of information. In order to see this, note that the system has ten parameters: three magnetic field components, three electric field components, three current components and the charge. There are eight independent relations connecting these quantities - the force free equation (2.1) has three constraints, the second order Maxwell's equation, (3.10), has three constraints as well and there are the two divergence relations involving the field components in (2.4) and (2.5). This leaves two independent quantities that are readily identified from the results of section 3 as the parallel (force free) current, $\mathbf{J}_\parallel$ and the displacement current orthogonal to the $\mathbf{k} - \mathbf{B}$ plane, δE_y .

4.1. Fast Wave Characteristics

The fast wave characteristics can be used to propagate finite changes in the displacement current orthogonal to the $\mathbf{k} - \mathbf{B}$ plane. Maxwell's equations can be used to find the corresponding induced magnetic field. By contrast, no information on the force-free current can be propagated along the fast characteristics. If one were to cover a region of spacetime with a mesh of short wavelength fast wave characteristics (i.e., the boundary of the spacetime region is a essentially a pure fast wave emitter and absorber), and time evolve the system

using the method of characteristics then one would find that the current and charge in the spacetime would remain unchanged for all time, in all frames.

There is a claim in Blandford (2001, 2002) that since the fast wave can transport a toroidal magnetic perturbation, “it carries information about the poloidal current.” This conjecture is inconsistent with the nature of the fast wave. The fast wave, like the light wave carries only displacement current. The time dependent version of Ampere’s law does not require a coupling between physical current and the magnetic field.

4.2. Alfven Wave Characteristics

By contrast, changes in the parallel current can be propagated along the Alfven wave characteristics. However, no changes in the displacement current orthogonal to the $\mathbf{k} - \mathbf{B}$ plane propagate along these characteristics. From equation (3.27) this current is essentially a pure charge moving along the magnetic field lines at the speed of light in the proper frame. The Alfven wave can thought of as a charge discontinuity or perturbation propagating at the speed of light along the magnetic field (and its associated parallel current) and the resulting changes in the electromagnetic field can be found from Maxwell’s equations.

Note that in the degenerate case of wave propagation parallel to the magnetic field, there is no $\mathbf{k} - \mathbf{B}$ plane. In the degenerate case, (3.24) implies that the Alfven wave carries no physical current, only a displacement current, δE_z .

4.2.1. The Force Free Plasma-Filled Waveguide

A very important application of the nature of the Alfven wave characteristics is a semi-infinite cylindrical waveguide threaded by a uniform axial magnetic field and filled with a force free plasma. A rotating conductive disk is suddenly attached to one end of the plasma-filled waveguide (see Chapter 2 of Punsly (2001) for pictures and complete details). The conductive disk behaves like a unipolar inductor or Faraday wheel. When the Faraday wheel is connected, a Poynting flux begins to be radiated down the plasma-filled waveguide. The Poynting flux is supported by a force free current. In the language of force free waves, the Faraday wheel has launched an axisymmetric force free Alfven step wave (see the Appendix B for details).

Note that by (3.26d), the Alfven wave needs to be oblique if it supports a field aligned current (i.e., a component of the propagation vector that is orthogonal to the field, $k_\perp$, is required). In the plasma-filled waveguide, the continuity of the tangential electric field at

the surface of the rotating disk and the degeneracy condition in the force free plasma imply that there is a poloidal electric field, $E_\perp$, that is launched by the Faraday wheel orthogonal to the poloidal magnetic field, B^P ,

$$E_\perp = -\frac{\Omega_D B^P \rho}{c}. \quad (4-1)$$

The coordinate, ρ , is the radial coordinate in a cylindrical coordinate system and the field line angular velocity, Ω_F is identical to the disk angular velocity, Ω_D . The radial variation of the electric field in (4.1) and in the oscillatory modes of (B1) is tantamount to a standing wave component of the propagation vector that is orthogonal to the magnetic field (note that the same wave polarization phenomenon occurs in vacuum electromagnetic waveguides). The step Alfvén wave in the cylinder (constructed in Appendix B) is a superposition (Fourier integral) of the oblique, axisymmetric Alfvén normal modes of the waveguide. The standing wave properties (obliquity) of the axisymmetric normal modes in the cylindrical waveguide are a consequence of the boundary conditions on the cylinder walls.

4.2.2. Pulsar Magnetospheres

One would expect a strong analogy between the plasma-filled waveguide and an idealized, force free, field aligned pulsar magnetosphere. The neutron star is the unipolar inductor, the magnetosphere is the plasma-filled waveguide and the wind is the Poynting flux in this analogy. Thus, one would expect that the pulsar wind is essentially an Alfvén wave radiated by the neutron star in this idealized limit.

Unfortunately, the causal nature of MHD in a relativistic magnetosphere was obfuscated by anecdotal comments in Blandford (2001) through an analysis in the force free approximation. If the force free analysis is not pathological then it must agree with the magnetically dominated limit of MHD. There is a claim that for axisymmetric modes, the Alfvén wave can only support variations in the poloidal magnetic field, not toroidal variations. However, this cannot be correct as the example of the plasma-filled waveguide shows by the explicit construction of the axisymmetric Alfvén modes that transport toroidal magnetic fields. The concept that a standing wave component of the propagation vector that is orthogonal to the magnetic field can exist in an axisymmetric mode (as in the plasma-filled waveguide) was not understood in Blandford (2001). The existence of such a wave vector polarization implies, with the aid of (3.30) that axisymmetric Alfvén modes that are capable of transporting toroidal field perturbations along the direction of poloidal magnetic field most certainly exist in a pulsar magnetosphere. This oversight allows for the claim that the force free analysis leads to an “opposite conclusion” to that of the MHD analysis. In reality, if one is prudent,

it can be shown as it has been done here that the two analyses agree, as they must.

5. Applications to Black Hole Magnetospheres

The force free approximation has been used in models of black hole magnetospheres in which spin energy is extracted electromagnetically from the black hole (Blandford and Znajek 1977; Thorne et al 1986). The force free wind systems in these treatments are characterized by two constants in each magnetic flux tube: the poloidal current (equivalently the toroidal magnetic field or angular momentum flux) and the field line angular velocity (this is quantity is equivalent to specifying the ratio of energy flux to angular momentum flux, the Goldreich-Julian charge density and the cross field potential).

The mathematical method of solution implements the force free condition at the horizon as a boundary condition that is used to determine the field aligned current density and the field line angular velocity in each flux tube. Since, by (3.17), the fast wave can propagate in any direction, at the speed of light, outgoing fast waves can propagate from anywhere outside of the event horizon (the fast critical surface for outgoing waves is the event horizon). By contrast, the "light cylinder" located within the ergosphere is the Alfvén critical surface, which no outgoing Alfvén wave can traverse (Blandford 2002; Punsly 2001). For rapidly rotating black holes, this surface is sufficiently far from the event horizon that the constraints imposed by the spacetime metric near the horizon are irrelevant. Thus, the use of the horizon boundary condition can only be justified from a causal perspective if the fast mode is involved in determining the wind constants. However, by (3.19), changes in the poloidal current or charge can not be transmitted along the fast wave characteristics in the black hole magnetosphere. Thus, the spacetime near event horizon can not affect changes in the wind parameters. The use of the event horizon as a causal wind boundary is not permitted in the force free approximation. Consequently, the mathematical method of solution is flawed. The same causal structure is known to exist in magnetically dominated perfect MHD (see Chapter 6 of Punsly (2001)). Going to the force free limit was conjectured in (Blandford 2001, 2002) to obviate any need for concern with this method of solution that were revealed in the full MHD analysis. This claim is clearly not supported by the results of this article.

The causal structure of the black hole magnetosphere in the force free limit has been shown, in this article to behave formally as the zero mass limit of MHD. As in perfect MHD, the wind parameters must be created outside of (farther out from the black hole than) the ergospheric Alfvén critical surface.

A further critique of the MHD analysis of black hole causality was that the Alfvén

wave really does not transport changes in current and charge, only oscillations (Blandford 2001). However, this statement is not an accurate depiction of the nature of propagating disturbances in MHD. It is well known that small amplitude MHD step waves obey the same dispersion relations as oscillatory waves. Alfven step waves and Alfven wave packets in general obey the same constitutive relations amongst field and dynamical quantities and propagate along the same characteristics as the oscillatory waves (Kantrowitz and Petschek 1966). This also follows by taking the appropriate Fourier integral to create a step wave from sinusoidal waves. The Alfven wave packets can transport finite changes in the charge density and field aligned currents in a force free magnetosphere. This is shown by explicit construction in Appendix B.

6. Discussion

There are three main reasons for not considering black hole magnetospheres that are force free **everywhere** to be physically realizable:

1. As mentioned in the Introduction, the physical plasma state near the event horizon is always inertially dominated. This physical state is exactly the opposite of the zero inertia limit that is inherent to the force free assumption.
2. From equation (2.7) there is no region within the force free magnetosphere where energy flux (Poynting flux) and angular momentum flux can be created. Energy and angular momentum flux (essentially, the poloidal current and Goldreich-Julian charge density) must be injected into the wind system at the boundaries. If the magnetosphere is force free everywhere then the energy flux (current and charge) must emerge from the horizon. However, the charge and current necessary to support the energy and angular momentum flux can not be propagated along the plasma wave characteristics emanating from the spacetime near the horizon as discussed in the last section. Thus, such a wind system is inherently acausal.
3. There are three MHD computer simulations of magnetically dominated black hole magnetospheres (the magnetic energy density exceeds the plasma energy density in the initial state) that have been published. All of them evolve to a state with strong cross-field (inertial, not force free) current flow in the ergosphere (Camenzind and Khanna 2000; Koide et al 2002; Semenov et al 2001). The inertial current flow initiates in regions far enough from the event horizon that it is not interpretable as a horizon surface current. The strong cross-field current is a necessary component for extracting

black hole energy in the theory of black hole gravitohydromagnetics (GHM) as well in Punsly (2001).

It should be noted that a wind system beyond (farther from the black hole than) the ergospheric dynamo region for the toroidal magnetic field (poloidal current) could in principle be well described by the force free approximation. From the nature of the force free characteristics, this dynamo region can exist only in the spacetime farther from the black hole than the ergospheric light cylinder (Alfven critical surface). The Poynting flux can propagate outward from the dynamo region to asymptotic infinity as an essentially force free wind. Such a wind system is described in detail in Chapter 9 of Punsly (2001).

I would like to thank Ferd Coroniti for his valuable and intelligent insights. I am also indebted to Serguei Komissarov for his useful comments regarding force free plasmas and propagating discontinuities.

A. First Order Curvature Effects - Hybrid Modes

In this appendix, we calculate the first order effects of field line curvature on the force free small amplitude plasma modes in a proper frame. One finds new hybrid modes as in MHD. The first order corrections to the short wavelength approximation can be found in the eikonal approximation. The sinusoidal solutions described in (3.2) are replaced by:

$$J^\mu = J_0^\mu + \delta J^\mu e^{i[\mathbf{k} \cdot d\mathbf{r} - \omega t]} , \quad (\text{A1})$$

where $\mathbf{k}$ is a slowly varying vector valued function. In the present circumstance, the unperturbed current no longer vanishes by assumption. However, by the force free condition, it flows parallel to the magnetic field in the proper frame. This can be expressed in terms of a slowly varying “radius of curvature”, R , as

$$\mathbf{J}_0 = cR^{-1}\mathbf{B}_0 . \quad (\text{A2})$$

The validity of the eikonal approximation in (A1) requires that:

$$|\mathbf{k}| R \gg 1 , \quad (\text{A3})$$

$$\frac{|\nabla k|}{k^2} \ll 1 . \quad (\text{A4})$$

The perturbed Maxwell’s equations and the perturbed degeneracy condition carryover from the sinusoidal case, (3.4) - (3.11), unchanged. The only perturbation equation that

changes is the force free condition, (3.2), that now becomes:

$$\mathbf{B}_0 \times \delta \mathbf{J} + \delta \mathbf{B} \times \mathbf{J}_0 = 0 . \quad (\text{A5})$$

The perturbed force free relation found from expanding the "x" and "z" components of (A5) is

$$\frac{4\pi i k^2 c^2}{c^2 k^2 - \omega^2} (kR)^{-1} \delta J_z = \delta J_y . \quad (\text{A6})$$

Note that by (A3), the large radius of curvature approximation, equation (A6) is approximately (3.12). The "y" component of the perturbed force free relation (A5) combined with (A6) yields the relation:

$$\frac{16\pi^2 k^4 c^4 (kR)^{-2} - [c^2 k^2 - \omega^2]^2}{[c^2 k^2 - \omega^2]^2} \frac{k_{\parallel}}{k} B_0 \delta J_z + \frac{k_{\perp}}{k} B_0 \delta J_x = 0 . \quad (\text{A7})$$

Note that in the limit of a large magnetic field radius of curvature that (A7) becomes (3.13). Thus, the complete set of perturbed equations reduce to the equations found in section 3 as the curvature of the field gets small.

The dispersion relation is found by combining (A7) with the perturbed degeneracy condition (3.14),

$$[\omega^2 - c^2 k^2] [\omega^2 - c^2 k_{\parallel}^2] = 16\pi^2 k^2 k_{\parallel}^2 c^4 (kR)^{-2} . \quad (\text{A8})$$

There are two roots of the dispersion relation.:

$$\omega^2 = c^2 k^2 + \frac{1}{2} c^2 \left[\sqrt{k_{\perp}^4 + 64\pi^2 k_{\parallel}^2 k^2 (kR)^{-2}} - k_{\perp}^2 \right] , \quad (\text{A9})$$

$$\omega^2 = c^2 k_{\parallel}^2 - \frac{1}{2} c^2 \left[\sqrt{k_{\perp}^4 + 64\pi^2 k_{\parallel}^2 k^2 (kR)^{-2}} - k_{\perp}^2 \right] . \quad (\text{A10})$$

Differentiating (A9) and (A10) yields the group velocities of the two modes. For parallel propagation, the group velocity is the speed of light up to terms that are $O[(kR)^{-2}]$. The parallel propagation assumption is invalidated at second order since the magnetic field is curved. In the limit that $(k_{\perp} R)^2 \gg 1$, one can obtain the following Taylor series expansions of the group velocity for oblique propagation:

$$\frac{d\omega}{dk} \approx \pm \frac{c}{\sqrt{1+16\pi^2 \cot^2 \theta (kR)^{-2}}} \approx \pm c [1 - 8\pi^2 (kR)^{-2} \cot^2 \theta] , \quad (\text{A11})$$

$$\frac{d\omega}{dk} \approx \pm \frac{c \cos \theta}{\sqrt{1-16\pi^2 \cot^2 \theta (kR)^{-2}}} \approx \pm c \cos \theta [1 + 8\pi^2 (kR)^{-2} \csc^2 \theta] . \quad (\text{A12})$$

Equations (A9) and (A11) approximate to (3.16) and (3.17) as the radius of curvature becomes infinite. Thus, these equations represent the fast hybrid mode. Similarly, (A10)

and (A12) become (3.22) and (3.23) in the short wavelength limit. These equations represent the hybrid Alfven wave. Notice that the fast hybrid mode propagates slower than the speed of light for oblique waves propagating across a curved background magnetic field. From (A12), the Alfven mode does not propagate perpendicular to the field. Thus, there is not Alfvenic properties for the fast mode to couple into this case. Thus, the hybrid fast mode still propagates at the speed of light for perpendicular propagation as indicated in (A11).

There are implications of this result to the primary focus of this article. Since the fast wave no longer propagates isotropically, it can now carry a physical current (not just a displacement current). Denote the displacement current as, J_D^y . Then (3.11a), (A6) and (A9) imply that the ratio of displacement current to physical current for oblique fast hybrid waves is:

$$\frac{J_D^y}{J_{\parallel}} \approx \frac{(kR)^3 \tan^4 \theta \sin \theta}{64\pi} . \quad (\text{A13})$$

Consequently, the ratio of displacement current to the physical current diverges in the short wavelength limit as in (3.19). Similarly, from the perturbed Gauss' law (3.7a), (3.11c) and (A9), one can find the ratio of displacement current to the charge density:

$$\frac{J_D^y}{\rho_e} \approx c \frac{(kR)^3 \tan^4 \theta \sin \theta}{64\pi} . \quad (\text{A14})$$

Comparing (A9) to (A13) and (A14), note that as the fast wave propagation speed slows down due to field line curvature, it carries more physical current and charge. It becomes increasingly more like the Alfven wave as the field line curvature increases. However, (A13) and (A14) show that in the eikonal approximation, the fast wave is still almost purely electrodynamic: a propagating displacement current with a minute physical current that is only $O[(kR)^{-3}]$ of the displacement current.

B. The Alfven Wave Packet in the Cylindrical Waveguide

In this appendix it is shown by explicit construction that a superposition of oscillatory Alfven waves (an Alfven wave packet) does indeed transport field aligned currents and charge, not just oscillatory perturbations. We rely on the magnetically dominated MHD treatment of the plasma-filled cylindrical waveguide found in Chapter 2 of Punsly (2001) and pass to the force-free limit. In order to save space, we do not reproduce the results of Chapter 2 here, but refer to them in detail and the reader is urged to consult that reference. In summary, a cylindrical waveguide filled with a tenuous plasma is placed inside of an infinitely long solenoid so that it is threaded by a strong axial magnetic field. The waveguide is semi-infinite in the sense that one end is terminated by a rotating conductive wheel. This rotating

conductor or Faraday wheel (also known as a homopolar generator or unipolar inductor) is an active element in the waveguide circuit that radiates plasma waves. In the first subsection, we construct the relevant axisymmetric, $m=0$, perfect MHD Alfven modes in the cylinder using the detailed results of Punsly (2001). In the next subsection, we compare and contrast a wave packet (a step wave) created from a linear superposition of the previously calculated oscillatory Alfven modes to the radiation emanating from the Faraday wheel (the nature of this radiation was derived previously in terms of a formal MHD wind solution in the waveguide in Punsly (2001)). Finally, it is noted that the depiction of the wind as a pure Alfven step wave becomes increasingly more accurate as one passes to the force-free limit.

B.1. Components of the Oscillatory Waves

This section constructs an oscillatory solution from Chapter 2 of Punsly (2001) for an axisymmetric Alfven wave, $m = 0$, in the plasma-filled waveguide that is consistent with the attachment of the Faraday wheel (in the realm of magnetically dominated perfect MHD).

B.1.1. The Electric Field

From equation (2.93b) of Punsly (2001), the radial electric field in an axisymmetric Alfven wave in a cylindrical waveguide is arbitrary and is determined by the boundaries (the field lines can shear relative to each other, hence these modes are often called “shear Alfven waves”). Considering the frozen-in condition within the Faraday wheel and the continuity of the tangential electric field at the Faraday wheel/plasma interface let us choose the radial electric field in the Alfven wave to be (in cylindrical coordinates):

$$E^\rho = -\frac{\Omega_D \rho}{c} B_0 e^{i(k_z z - \omega t)} . \quad (\text{B1})$$

The other electric field components are given in Punsly (2001) by (2.86c) and (2.93a) as

$$E^\phi = 0 , \quad (\text{B2})$$

$$E^z = 0 . \quad (\text{B3})$$

B.1.2. The Currents

From equations (B1)-(B3) and Gauss’ law, the charge density is;

$$\rho_e = -\frac{\Omega_D B_0}{2\pi c} e^{i(k_z z - \omega t)} . \quad (\text{B4})$$

From the Alfven wave dispersion relation, (2.82) of Punsly (2001), and the frozen-in form of the momentum equation, (2.86a) of Punsly (2001), along with equation(A1), the cross-field current density is

$$J^\rho = i \frac{v_I k_z \Omega_D \rho}{4\pi U_A^2 c} B_0 e^{i(k_z z - \omega t)} , \quad (\text{B5})$$

where v_I is the intermediate three speed and U_A is the pure Alfven speed. In the force free limit, the pure Alfven speed diverges as the intermediate speed approaches the speed of light. From the law of current conservation,

$$\frac{\partial \rho_e}{\partial t} + \nabla \cdot \mathbf{J} = 0 , \quad (\text{B6})$$

combined with (B4) and (B5) yields the axial current density:

$$J^z = \frac{\rho_e}{v_I} c^2 = - \frac{\Omega_D B_0 c}{2\pi v_I} e^{i(k_z z - \omega t)} . \quad (\text{B7})$$

B.1.3. The Magnetic Field

Using the value of the axial current in (B7) in Ampere's law yields the toroidal magnetic field strength:

$$B^\phi = - \frac{B_0 \Omega_D \rho}{v_I} e^{i(k_z z - \omega t)} . \quad (\text{B8})$$

The other components are given by (2.92a) and (2.96) of Punsly (2001):

$$B^z = 0 , \quad (\text{B9})$$

$$B^\rho = 0 . \quad (\text{B10})$$

B.2. Making Wave Packets

We can construct wave packets of the oscillatory solutions above by taking Fourier integrals. We are interested in the step wave. For the Alfven wave, one can use the dispersion relation (2.82) of Punsly (2001) in the cylinder to write,

$$e^{i(k_z z - \omega t)} = e^{-ik_z(v_I t - z)} . \quad (\text{B11})$$

The step function is given by,

$$\Theta(v_I t - z) = - \frac{1}{2\pi i} \int_{-\infty}^{+\infty} \frac{e^{-ik_z(v_I t - z)} dk_z}{k_z + i\epsilon} . \quad (\text{B12})$$

We construct a wave packet of Alfven waves, Ψ , with a spectral amplitude, $A(k_z)$,

$$\Psi = \int_{-\infty}^{+\infty} A(k_z) \psi(k_z) dk_z , \quad (\text{B13})$$

$$A(k_z) = -\frac{1}{(2\pi i)(k_z + i\epsilon)} , \quad (\text{B14})$$

where $\psi(k_z)$ is one of the oscillatory wave function components in equations (B1)-(B10) and ϵ is an arbitrarily small positive number in the usual sense. The wave packet of oscillatory solutions with field and current components given by (B1)-(B10) and spectral amplitude given by (B14) has the following field and current distributions (for the sake of demonstrating equivalence, the corresponding equation number from Punsly (2001) that were derived by the MHD wind calculation in the cylinder is placed on the right hand side of the equals sign in these relations):

$$E^\rho = -\frac{\Omega_D \rho}{c} B_0 \Theta(v_I t - z) = (2.98) , \quad (\text{B15})$$

$$E^\phi = 0 , \quad (\text{B16})$$

$$E^z = 0 , \quad (\text{B17})$$

$$\rho_e = -\frac{\Omega_D B_0}{2\pi c} \Theta(v_I t - z) = (2.99) , \quad (\text{B18})$$

$$J^\rho = \frac{v_I \Omega_D \rho}{4\pi U_A^2 c} B_0 \delta(v_I t - z) , \quad (\text{B19})$$

$$J^z = \frac{\rho_e}{v_I} c^2 = -\frac{\Omega_D B_0 c}{2\pi v_I} \Theta(v_I t - z) = (2.119) , \quad (\text{B20})$$

$$B^\phi = -\frac{B_0 \Omega_D \rho}{v_I} \Theta(v_I t - z) = (2.118) , \quad (\text{B21})$$

$$B^z = 0 , \quad (\text{B22})$$

$$B^\rho = 0 . \quad (\text{B23})$$

These quantities agree with the downstream state of the plasma that was found near the flow front in Punsly (2001) in the limit that $v_z = v_I$. Notice that J^ρ vanishes downstream of the wavefront and only has a surface component on the wavefront as it does in equation (2.110) of Punsly (2001).

B.3. Conclusion

From equations (B18) and (B20), the Alfven wave packet actually transports charge and field aligned current.

The solution is not exact. There are errors associated with the small inertial terms in the magnetically dominated limit. At the wavefront, the relativistic MHD shock equations do not solve exactly. These equations are the conservation of the axial components of the stress energy tensor across the discontinuity in the frame of the wavefront. Pressure balance (magnetic pressure from the toroidal magnetic field in the frame of the wavefront) can not be achieved because the Alfven wave, unlike the fast wave, has no compressive properties. The errors in the stress-energy balance can be found in the frame of the propagating discontinuity by using (B15) and (B21) to compute the proper toroidal magnetic field. The errors in the shock equations are on the order of $(\Omega_D \rho U_A^{-1})^2$. This is biquadratic in two small quantities: $\Omega_D \rho / c \ll 1$ by construction and by the magnetically dominated condition, $c U_A^{-1} \ll 1$. The violation of the shock relations are therefore a very small second order effect in the magnetically dominated limit. Notice that the errors vanish completely in the force-free limit as the Alfven three speed approaches the speed of light. In magnetically dominated perfect MHD, these small errors are accounted for by a fast switch-on shock that creates a B^ρ and an E^ϕ downstream of the shock front. This is an MHD precursor, an infinitesimal distance upstream, to the Alfven rotational discontinuity at the terminus of the Alfven step wave. As demonstrated by explicit construction above, it is the Alfven rotational discontinuity that imprints the charge and field aligned current on the waveguide plasma. The interpretation of the waveguide wind solution as an Alfven wave is exact in the force-free limit and is extremely accurate to first order in the magnetically dominated MHD limit. However, in relativistic MHD it is important to note that both fast and Alfven modes are needed to solve this type of “Riemann problem.”

It was an oversight in Punsly (2001) that nonlinear fast waves such as an abrupt discontinuity can in principle transport charge and current. However, in practice, due to the limited polarization properties of such waves (the fast discontinuity can only affect changes in the magnetic field in the plane spanned by the normal to the wavefront and the magnetic field upstream) unless the magnetic field is perfectly homogenous (not the case in a magnetosphere) they must be accompanied by an Alfven rotational discontinuity in a magnetosphere in order for the discontinuities to be solutions of the relativistic shock equations (Kantrowitz and Petschek 1966). The important point is that in general both modes are needed. The discussions of Punsly (2001) were designed to show that the Alfven wave is always required in MHD, so that the fast wave cannot determine the wind constants alone - and this conclusion is unchanged and hence the MHD causality arguments are unchanged. In the force-free limit, the situation is more extreme. Since the fast wave is linear in the force free limit (the fast speed is always c), all wave packets constructed from the complete set of oscillatory fast modes will also carry no current or charge in any frame (see section 3.1). Thus, the Alfven wave is responsible for setting the wind constants in a force free black hole magnetosphere.

This discussion corroborates the physical accuracy of the assumptions used in the one dimensional string analysis of flux tubes in Semenov et al (2001). They actually ignore the fast mode completely (because as they say, the fast wave is well known to primarily provide only pressure changes in an MHD flux tube) and only consider the Alfvén mode. This seems like a very reasonable assumption based on the analysis provided here.

REFERENCES

- Blandford, R., Prog. Theor. Phys. Supp. 2001, **147**, 182
- Blandford, R. 2002, astro-ph0202265 to appear in “Lighthouses of the Universe,” eds. Gilfanov, M. et al.(Springer:Berlin)
- Blandford, R. and Znajek, R. 1977, MNRAS. **179**, 433
- Camenzind, M. and Khanna, R. 1999, Nuovo Cimento **B115**, 815
- Hawking, S. 1973, In: *Black Holes, Les Houches 1973* ed. by B. DeWitt and C. DeWitt (Gordon and Breach, New York)
- Kantrowicz, A.R. and Petschek, H.E. 1966, In: *Plasma Physics in Theory and Application*. ed. by W.B. Kunkel (McGraw-Hill, New York) p. 148
- Koide, S., Shibata, K., Kudoh, T. and Meier, D. 2002, Science **295**, 1688
- Lanzerotti, L.J. and Southwood, D.J. 1979, In: *Solar System Plasma Physics* vol. III ed. by E.N. Parker, C.F. Kennel, L.J. Lanzerotti (North Holland, Amsterdam)
- Thorne, K., Price, R. and Macdonald, D. 1986, *Black Holes: The Membrane Paradigm*. (Yale University Press, New Haven)
- Punsly, B. 2001, *Black Hole Gravitohydromagnetics* (Springer-Verlag, New York)
- Semenov, V. 2002, et al, Physica Scripta **65**, 13
- Uchida, T. 1997, MNRAS. **291**, 125